## **BMC Pharmacology**



Poster presentation

**Open Access** 

# Internalization and degradation of natriuretic peptide receptor-A is stimulated by ligand binding

Darcy R Flora\*1, Sean D Conner<sup>2</sup> and Lincoln R Potter<sup>1,3</sup>

Address: <sup>1</sup>Department of Pharmacology, University of Minnesota, Minneapolis, MN, USA, <sup>2</sup>Department of Genetics, Cell Biology and Development, University of Minnesota, Minneapolis, MN, USA and <sup>3</sup>Department of Biochemistry, Molecular Biology and Biophysics, University of Minnesota, Minneapolis, MN, USA

Email: Darcy R Flora\* - flora097@umn.edu

\* Corresponding author

from 4th International Conference of cGMP Generators, Effectors and Therapeutic Implications Regensburg, Germany. 19–21 June 2009

Published: 11 August 2009

BMC Pharmacology 2009, 9(Suppl 1):P14 doi:10.1186/1471-2210-9-S1-P14

This abstract is available from: http://www.biomedcentral.com/1471-2210/9/S1/P14

© 2009 Flora et al; licensee BioMed Central Ltd.

### **Background**

Natriuretic peptide receptor-A (NPR-A) is a transmembrane receptor guanylyl cyclase that binds and mediates the effects of atrial and B-type natriuretic peptides (ANP/BNP). Internalization and ligand-dependent degradation of NPR-A is controversial, in part due to the use of ligand binding studies to predict the cellular location of the receptor. Here, we used a more direct sequential immuno-precipitation-western blot assay to demonstrate that long-term ANP exposure increases NPR-A degradation in primary, immortalized, and transfected cells.

#### **Results**

A separate novel extracellular epitope antibody-binding assay indicated that NPR-A is internalized under basal conditions and that this rate is increased about two-fold by ANP exposure. siRNA knock down of clathrin and dominant negative inhibition of dynamin failed to inhibit ANP-dependent NPR-A degradation, whereas dominant negative dynamin expression reduced the rate of NPR-A internalization about 40%.

#### **Conclusion**

These data indicate that NPR-A is basally internalized by a dynamin-dependent pathway and that prolonged ANP exposure stimulates both NPR-A internalization and degradation.

#### Acknowledgements

D.R.F. was supported by an award from the American Heart Association (0815607G).