

Meeting abstract

Open Access

## The gut-mood axis: a novel role of the gut hormone peptide YY on emotional-affective behaviour in mice

Evelin Painsipp\*<sup>1</sup>, Herbert Herzog<sup>2</sup> and Peter Holzer<sup>1</sup>

Address: <sup>1</sup>Institute of Experimental and Clinical Pharmacology, Medical University of Graz, 8010 Graz, Austria and <sup>2</sup>Neurobiology Research Program, Garvan Institute of Medical Research, Sydney, NSW 2010, Australia

Email: Evelin Painsipp\* - evelin.painsipp@medunigraz.at

\* Corresponding author

from 15th Scientific Symposium of the Austrian Pharmacological Society (APHAR) Joint meeting with the Hungarian Society of Experimental and Clinical Pharmacology (MFT) and the Slovenian Pharmacological Society (SDF) Graz, Austria. 19-21 November 2009

Published: 12 November 2009

BMC Pharmacology 2009, 9(Suppl 2):A13 doi:10.1186/1471-2210-9-S2-A13

This abstract is available from: <http://www.biomedcentral.com/1471-2210/9/S2/A13>

© 2009 Painsipp et al; licensee BioMed Central Ltd.

### Background

Peptide YY (PYY) and neuropeptide Y (NPY) are members of the PP-fold peptide family. PYY is expressed by endocrine cells of the gut, whereas NPY occurs in enteric, sensory, cerebral and autonomic neurons. While PYY is involved in the regulation of gut function and satiety, NPY is known to contribute to the regulation of emotional-affective behaviour, cognition, seizure activity, pain and satiety. Since we have previously found that the NPY system plays a role in the gut-brain axis, we explored in which way knockout of PYY and PYY plus NPY alters the emotional-affective behaviour in mice and how this phenotype is altered by experimental colitis.

### Methods

Male wildtype (WT), PYY (PYY<sup>-/-</sup>) and PYY plus NPY (PYY+NPY<sup>-/-</sup>) knockout mice, all with a mixed C57BL/6:129/SvJ (1:1) background, were used. Mild colitis was induced by adding dextran sulphate sodium (DSS, 2%) to the drinking water for 7 days. Inflammation was assessed by the colonic myeloperoxidase content, anxiety-related behaviour was evaluated with the elevated plus maze (EPM) and open field (OF) tests, and depression-like behaviour was estimated with the forced swim test (FST).

### Results

When the animals were phenotyped in the absence of colitis, anxiety-like behaviour in the EPM (reduction of open arm time) was increased in both PYY<sup>-/-</sup> and PYY+NPY<sup>-/-</sup>

mice, whereas locomotion remained unaltered. The results obtained in the OF test were largely similar. The depression-like behaviour (immobility) in the FST was markedly enhanced in both PYY<sup>-/-</sup> and PYY+NPY<sup>-/-</sup> mice. DSS-induced colitis was associated with an increase in the colonic myeloperoxidase content. On the behavioural level, colitis had genotype-dependent effects on emotional-affective behaviour. Most conspicuous was that colitis increased anxiety primarily in WT mice so that in the EPM test WT mice spent less time on the open arms than the knockout animals. Similar changes were observed in the OF test. In contrast, colitis did not alter depression-like behaviour in WT and PYY+NPY<sup>-/-</sup> mice but reduced immobility in PYY<sup>-/-</sup> mice.

### Conclusion

These data show for the first time that the gut hormone PYY has a significant impact on emotional-affective behaviour, because its deletion enhances anxiety- and depression-related behaviour. Colitis enforces anxiety-like behaviour only in the presence of PYY, whereas in the absence of PYY depression is less pronounced in animals with colitis. These data attest to an important role of PYY in the gut-mood axis.