

Poster presentation

Open Access

Blockade of CNG channels abrogates urethral relaxation induced by soluble guanylate cyclase activation

Domingo Triguero*, Maria Sancho, Marta García-Flores and Ángeles García-Pascual

Address: Department of Physiology, Veterinary School, Complutense University, 28040 Madrid, Spain

Email: Domingo Triguero* - dtriguer@vet.ucm.es

* Corresponding author

from 4th International Conference of cGMP Generators, Effectors and Therapeutic Implications Regensburg, Germany. 19–21 June 2009

Published: 11 August 2009

BMC Pharmacology 2009, 9(Suppl 1):P69 doi:10.1186/1471-2210-9-S1-P69

This abstract is available from: <http://www.biomedcentral.com/1471-2210/9/S1/P69>

© 2009 Triguero et al; licensee BioMed Central Ltd.

In the present study, we have characterized the presence and distribution of cGMP-gated cationic channels (CNG) in the rat urethra as well as its putative role in the mediation of the nitrenergic relaxation. Previous studies have shown the inhibition of the sheep urethral nitrenergic relaxations by the CNG's inhibitor L-cis-diltiazem [1]. Also in the rat urethra, L-cis-diltiazem (50 μ M) inhibited nitrenergic relaxations elicited by electrical field stimulation (EFS) of arginine-vasopressin (AVP)-precontracted urethral preparations (Figure 1A).

This effect was stereoselective since the isomer D-cis-diltiazem did not have any effect on urethral relaxations. Similarly, L-cis-diltiazem (50 μ M) showed a complete blockade of the relaxation induced by the addition of YC-1 (50 μ M), an specific activator of the soluble guanylate cyclase (sGC) (Figure 1B), thus reinforcing that L-cis-diltiazem would be acting as specific inhibitor of CNG channels in this tissue.

The presence of these channels in urethral tissue was tested by carrying out conventional RT-PCR studies using retina as a positive control. A band of the predicted size showing after sequencing, a 99% identity with the rod-type CNGA1, together to a second band with a similar size to that of the CNGB1 fragment, were obtained. These results suggest that they are heteromeric retina-like CNG channels.

Immunofluorescence studies were performed to analyze the distribution of CNG immunoreactivity (-ir) in sections of the urethral wall. As can be seen in Figure 2, a strong CNG-ir was present in a subpopulation of vimentin-ir cells known as Interstitial Cells of Cajal (ICC), while




Figure 1
The presence of L-cis-diltiazem (50 μ M) rapidly and effectively inhibited relaxation in arginine-vasopressin (AVP)-precontracted urethral preparations induced by both A) EFS (0.8 ms, 10 Hz, 5 s at 2 min-intervals) and B) guanylate cyclase activation by YC-1 (50 μ M).




Figure 2
Section of rat urethra showing immunofluorescence for CNG1 channels (green) and vimentin (red) and its co-localization in an intramuscular Interstitial cell [nuclei are stained with DAPI (blue)]. Scale bar: 20 μ m.

a fainter labeling was observed in smooth muscle cells (identified by actin-ir). CNG-ir was never observed in neural structures, nerve terminals, nerve trunks or ganglia. This pattern of CNG labeling and co-localization is similar to that described for the cGMP-ir found in the urethra upon functional nitrenergic stimulation [2]. We suggest that CNG channels could be an essential ionic mechanism that would link activation of the NOS-GC pathway to relaxation. In addition, these results point to ICC as key elements in the mediation of the nitrenergic urethral relaxation, although its precise role remains to be elucidated.

Acknowledgements

This work was supported by a grant (BFU2006-15135-C02-01) from the Ministerio de Educación y Ciencia of Spain.

References

1. Triguero D, González M, García-Pascual A, Costa G: **Atypical relaxation by scorpion venom in the lamb urethral smooth muscle involves both NO-dependent and -independent responses.** *Naunyn-Schmiedeberg's Arch Pharmacol* 2003, **361**:151-159.
2. García-Pascual A, Sancho M, Costa G, Triguero D: **Interstitial cells of Cajal in the urethra are cGMP-mediated targets of nitrenergic neurotransmission.** *Am J Physiol Renal Physiol* 2008, **295**:F971-F983.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

